
## **Changing Crabs**





Monitoring the arrival of a new species with citizen science

**Alderney** 

Wildlife Trust

# Turn back the clock to 2017

Trumps first inauguration as president

 Hull is the UK city of culture

and Despacito is top of the charts





## Turn back the clock to 2017

### But in Alderney, we are focused on crabs

• Fishers have noticed declining catches and increased disease







## Preliminary surveys

Dipping our toe into the proverbial rock pool



- Is the new species a potential problem?
- Are there signs of disease in the intertidal zone?






Edible crab ('AKA' Chancre)

Cancer pagurus







Green shore crab

Carcinus maenas







Velvet swimming crab

Necora puber







Isle of Man

A range of other infrequent species e.g. this hairy crab

Pilumnus sp.



# Furrowed crab Xantho hydrophilus



And two 'new' species, driven by climate change




Risso's crab

Xantho pilipes

**Alderney** Wildlife Trust





## The Big Three



Edible crab ('AKA' Chancre)

Cancer pagurus





Green shore crab

Carcinus maenas





Furrowed crab

Xantho hydrophilus



### Preliminary survey results









Year

n. Surveys

Green shore Edible crab crab

**Furrowed** crab





# Preliminary survey results








Year	n. Surveys	Green shore crab	Edible crab	Furrowed crab
2017- 2018	2	15	8	0

### Partially-standardised surveys

2020	4	2	24	24
2021	11	1	41	70





# Preliminary survey results








Year	n. Surveys	Green shore crab	Edible crab	Furrowed crab
2017- 2018	2	15	8	0

#### Partially-standardised surveys


2020	4	2	24	24
2021	11	1	41	70
















## The preliminary surveys had some significant drawbacks

Without standardized surveys we cannot tell what is really happening

But without resources how can we conduct standardized surveys!?








## Crab Surveys

We designed a new method for our 'Marine Group' to carry out

Aims to record
abundance of all
species and
biometrics of
arget species.
















## Mean # crabs per rock 2022-2025




Mean # crabs per rock 2022-2025

IIEM 25




Wildlife Trust


### Shore crab locations




















## With freshwater input


IIEM 25




## Without freshwater input (Longis Removed)



## Without freshwater input (Longis Removed)





IIEM 25




Edible crab ('AKA' Chancre)


Cancer pagurus












Furrowed crab


Xantho hydrophilus








## Results - Size Profile



## Results - Size Profile






Juvenile shore crabs spend most of their time in the intertidal zone but adults migrate into the subtidal<sup>2</sup>




## Results – Size Profile



**Green Shore Crab** 

Normal distribution indicates all adult age cohorts are in the intertidal zone.




**Alderney**Wildlife Trust



This is supported by high proportion of female furrowed crabs having eggs in summer









This is supported by high proportion of

female furrowed crabs having eggs in

summer










#### Results - Size Profile








Edible Crab

## Sex ratio of edible crab found in Alderney since 2022










## Sex ratio of edible crab found in Alderney since 2022






In Anglesey sex ratio of 1: **1.77** In Cornwall sex ratio of 1: **0.6** In Alderney, sex ratio of 1: **13.6** 





#### Results - Size Profile








- •
- M
- (






- Furrowed crab have become the most abundant on the rocky shore in just a few years
- This is a climate change driven shift
- Changes such as these are only going to continue as sea temperatures warm





- Green shore crab appear to be restricted to brackish areas
- Could this be due to competition?
- Changes in rainfall are also altering our freshwater inputs
- With no estuaries or other large brackish areas, shore crabs may be vulnerable in small islands






 Edible crab are infrequently encountered, but no clear declines (SINCE 2022)

 Good sign, given concerns over disease and warming temperatures affecting their life cycle


However, we find very few.... and








- Alarming sex ratio in edible crabs
- Temperature? Behaviour?
   Sampling error?
- And what is the ratio in adults?














Desk based review







 Refine method and investigate targeted projects, e.g. MSc

• Other citizen science – *crab* most wanted?











Subtidal Monitoring




 Reduce carbon in the atmosphere






- Reduce carbon in the atmosphere
- Consider conservation action











Wildlife Trust

- 1. Silva, A. (2008) *Predation by crabs on rocky shores in north-east Atlantic.* Thesis. University of Plymouth. Available at: <a href="http://dx.doi.org/10.24382/3535">http://dx.doi.org/10.24382/3535</a>
- 2. Rowley, A.F. & Coates, C.J. (2023) Shell disease syndromes of decapod crustaceans. *Environmental Microbiology*, 25(5), 931–947. Available from: <a href="https://doi.org/10.1111/1462-2920.16344">https://doi.org/10.1111/1462-2920.16344</a>
- 3. Naylor, E. "Seasonal Changes in a Population of Carcinus Maenas (L.) in the Littoral Zone." *Journal of Animal Ecology*, vol. 31, no. 3, 1962, pp. 601–09. *JSTOR*, https://doi.org/10.2307/2055. Accessed 5 Sept. 2025.
- 4. Heraghty, N. (2013). Investigating the abundance, distribution and habitat use of juvenile Cancer pagurus (L.) of the intertidal zone around Anglesey and Llŷn Peninsula, North Wales (UK). MSc thesis, Bangor University, Fisheries & Conservation report No. 29, Pp.75
- 5. Jenkin, A., Trundle, C., Sturgeon, S., Daniels, C. and Street, K. 2023. Intertidal Edible Crab (Cancer pagurus) Survey 2022. Cornwall Inshore Fisheries and Conservation Authority (Cornwall IFCA), Hayle.
- 6. Weiss, Monika & Thatje, Sven & Heilmayer, Olaf & Anger, Klaus & Brey, Thomas & Keller, Martina. (2009). Influence of temperature on the larval development of the edible crab, Cancer pagurus. Journal of the Marine Biological Association of the United Kingdom. 89. 753 759. 10.1017/S0025315409003269.



## Thank You for Listening!

#### **Key findings**

- 1. Furrowed crabs are now very abundant
- 2. Shore crabs only abundant in brackish areas
- 3. Edible crabs are uncommon
- 4. Sex ratio of 1:14 female: male edible crabs

#### What's next?

- Assess priorities and adapt survey
- Desk review & MSc?
- Subtidal data
- Consider campaigning to protect key areas

#### **Acknowledgements**

Alderney Marine team and volunteers, in particular: Dr Mel Broardhurst-Allen, Lou, Bob, Mark, Diana, Esme, Nigel, and many others.



Alex Purdie

<u>awte@alderneywildlife.or</u>

g

