



The Isle of Man Shark Tagging Programme

End of Year Report 2025



Written for:

The Department of Environment, Food and Agriculture (DEFA)

Written by Tom Morgan and Dr Lara Howe

Introduction

The Small Shark Tagging Programme in the Isle of Man has been operating since May 2013, with the Manx Wildlife Trust (MWT) working in partnership with the Department of Environment, Food and Agriculture (DEFA) to collect data. Sharks, skates and rays are currently subject to multiple threats from fisheries and harvest, including small-scale subsistence fishing, large-scale harvesting and unintentional bycatch. These species are therefore protected in many jurisdictions. However, little is known about the distribution, movement or population sizes of these cryptic species in Manx waters. The Small Shark Tagging Programme aims to work with local anglers to tag small sharks and rays with identification tags or streamers, on a catch and release basis. It is hoped the data will provide much needed information on the distribution and numbers of these small shark populations. Going forward, this fundamental understanding is crucial in providing effective and evidence-based data for the future management of these species and the best ways to protect them. The present report is a continuation of this programme, summarising the findings from this year.

Since the Small Shark Tagging Programme's inception, 811 individual elasmobranchs, made up of four species, have been caught and 809 of those tagged in Manx waters. These species include bull huss (*Scyliorhinus stellaris*), spurdog (*Squalus acanthias*), thornback ray (*Raja clavata*) and tope (*Galeorhinus galeus*). A key success of the programme is the recapture of 12 tagged tope, a further eight tope were caught in Manx waters tagged in other programmes. This highlights the mobile nature of these sharks and the need for international cooperation on their protection.

Due to the COVID-19 pandemic, the Small Shark Tagging Programme was unable to tag any small sharks or complete further training during 2020. The project resumed in 2021 and has continued in the subsequent years with further training provided. This year, 8 anglers have been trained, bringing the total number of anglers trained to 111.

The Scottish Shark Tagging Programme greatly contributed to the inception of this programme and showed what can be achieved through citizen science. Although the Scottish Shark Tagging Programme disbanded in 2018, the programme was a great source of knowledge and resources to the MWT programme. This included the deployment of two officers who trained Manx local anglers in 2013 (funded by DEFA), the design of a project logo, and the annual provision of tags/tagging equipment. Then again in 2017, an officer joined MWT for further training and support. In addition, they also increased public awareness highlighting the need for shark protection, the importance of sea angler's conservation efforts, and contributed to over 26 species of sharks, skates and rays being protected in Scottish waters.

Project Aims:

- Promote public awareness on the importance of small shark species and the need for their protection.
- Engage with local anglers to undertake tagging and record subsequent recaptures.
- Utilise the data collected to determine the abundance and distribution of Manx small shark populations.
- Examine local threats to small shark species to inform management plans and conservation activities.

Species Overview

Bull huss

Bull huss are globally classified as 'Vulnerable' but listed as 'Near Threatened' in Europe by the IUCN Red List, with an overall decreasing population trend (Figure 1; Ellis *et al.*, 2015a; Finucci, Derrick and Pacoureaux, 2021). Falling population trends are due to continuously declining numbers of mature individuals and severely fragmented populations (Finucci, Derrick and Pacoureaux, 2021). Bull huss experience a high level of exploitation across the species known range, with an overall population reduction of 30% - 49% over the last 48 years (Sherley *et al.*, 2020, Winker *et al.*, 2020). Contrary to this, standardised catch-per-unit-effort (CPUE) data in the Irish Sea and Bristol Channel showed an annual rate of increase of 4.7%. Bull huss is thereby considered locally abundant with a regionally increasing population around the British Isles (ICES-WGEF, 2019).

Females lay eggs two at a time between March and October, taking between 7-12 months to hatch. Sexual maturity is attained for males at 77 cm total length (TL), and females at 79 cm TL, corresponding to an age of four years if hatchling growth rates remain constant (Capapé *et al.*, 2006). This species has a lifespan of approximately 17 - 19 years (Rodríguez-Cabello *et al.*, 2005).




Figure 1 The IUCN Red List of Threatened Species categories.

Spurdog

Spurdog are classified as 'Vulnerable' globally by the IUCN Red List (Figure 1; Fordham *et al.*, 2016; Walker *et al.*, 2006), but 'Endangered' in Europe (Ellis *et al.*, 2015b). This assessment is based on a continuing decline in the number of mature individuals and severely fragmented populations. Population genetics has revealed little to no genetic mixing between Northern and Southern Hemisphere populations. Even where stocks have overlapping geographic range, mixing remains limited (Veríssimo *et al.*, 2010). Within European waters, it is suspected there are three distinct subpopulations (Northeast Atlantic, Mediterranean Sea and Black Sea; Veríssimo *et al.*, 2010). Spurdog are highly migratory, travelling in large, dense aggregations, segregated by size and sex. This aggregating behaviour makes CPUE data an unreliable indicator of population status, as a high CPUE may be maintained even when populations are severely depleted (Ellis *et al.*, 2015b).

Spurdog breed every other year (Holden and Meadows, 1962; Sosinski, 1978; Fahy, 1989), likely mating offshore in winter (Castro, 1983; Compagno, 1984), between October and February (Jones and Ugland, 2001). The central Irish Sea has been suggested as a possible key mating site (Dureuil, 2013). Females sexually mature at 74 - 92.5 cm TL and males mature at 57.5 - 64 cm TL (Henderson *et al.*, 2002). The maximum age is at least 40 years (Fahy, 1989), and fecundity increases with size (Ellis and Keable, 2008).

Thornback ray

Thornback ray are classified as 'Near Threatened', with population numbers considered stable in European waters (Ellis, 2016; Ellis *et al.*, 2016). Thornback ray is widespread and one of the more abundant elasmobranchs in the inner continental shelf seas of Europe. Scientific trawl surveys in northern European waters show that relative abundance has been stable or increasing in recent years, following long-term historical declines (McHugh *et al.*, 2011). However, catch analysis from 2000 to 2006 in the North Sea showed 38% of thornback ray were caught prior to reaching sexual maturity (64 cm/~ five years of age; Wiegand *et al.*, 2011). If current fishing patterns continue, it could result in a projected population decline of 90% within 30 years. Therefore, although research indicates that catch rates have increased in core parts of the geographic distribution, this is a recent increase following a longer-term decline.

Kadri *et al.*, (2014) found average age to be at least 15 years. The estimated size at 50% maturity for females have been estimated at 45 cm disc width (DW) and 77 cm TL (eight years), and for males at 42 cm DW and 66.6 cm TL (seven years; Walker, 1998; McCully *et al.*, 2012). This species first spawns in its fifth year (Ryland and Ajayi 1984). The fecundity in British waters has been estimated at between 100 - 140 eggs per year (Holden, 1975; Ryland and Ajayi, 1984). It may not be a continuous spawner and release around 35 eggs over four clutch episodes (Serra-Pereira *et al.*, 2011). The nursery areas used are coastal areas (e.g., the Wash and Thames estuary in the UK).

Tope

Tope are classified as 'Critically Endangered' globally but listed as 'Vulnerable' in Europe by the IUCN Red List, with an overall decreasing population trend (Figure 1; Walker *et al.*, 2020; McCully, Dureuil and Farrell, 2015). Declining population trends have been accredited to severely fragment populations and a continuing decline of mature individuals (McCully, Dureuil and Farrell, 2015). There is believed to be a single stock of tope in the Northeast Atlantic, extending from southern Norway and Scotland, southwards to the coast of northwest Africa and the Mediterranean Sea (ICES, 2012). Tope landings throughout the Northeast Atlantic region have decreased by 83% in the past twenty years (ICES, 2011). Landings off west Scotland, the Celtic Seas and the English Channel are relatively high, but highly variable (ICES, 2012). Greater abundances of mature individuals have also been found off the northern Irish coast, the southern Irish Sea and the east coast of England (Dureuil, 2013).

This species typically occurs in schools, partially segregated by size and sex (Walker *et al.*, 2008). There is regional variation in size at maturity, in the Northeast Atlantic males mature at 121 cm TL and females mature at 155 cm TL (Dureuil, 2013). Female age-at-maturity varies from 10 - 15 years (average 12.5 years) and maximum age is estimated as 40 years (tag returns suggest a possible maximum age of 60 years; Olsen, 1954; Francis and Mulligan, 1998; Walker, 1999). Tope reproductive cycles vary regionally from annual to triennial, although studies with intensive sampling indicate triennial cycles (Walker *et al.*, 2006; Ebert, Fowler and Compagno, 2013). Average litter sizes are between 20 to 35 pups, with litter sizes increasing in larger females (Ebert, 2003). Shallow, protected bays and estuaries serve as pupping and nursery area where young remain for up to two years (Walker *et al.*, 2006; Bovcon, 2018; McMillan *et al.*, 2018).

Methodology

The project is advertised locally and experienced anglers targeting small sharks are invited to partake in the programme. In 2025, 8 anglers were trained to administer tags. Since the beginning of the programme, 111 anglers have received small shark tagging training.

All trained anglers were given a minimum tagging size crib sheet, recording cards and tagging equipment (Appendix 1 and 2). Prior to tag application, the condition of each shark is visually assessed to ensure normal appearance and minimum tagging size. Any injured or otherwise abnormally appearing sharks, or those below the minimum tagging size, are rejected from the tagging pool. Information is recorded on the including: species, location, date, length, girth, sex and condition. Tagging equipment consists of a micro gun, with ten micro-tags for tagging smaller sharks. Tag equipment is replaced in small quantities when required, depending on angler's likelihood of being able to fish. An external tag with imprinted unique identification numbers is applied to each fish, which is also recorded on the recording card. Anglers are able to email tagging information directly to the MWT. The micro-tags are inserted at a 45° angle, then the trigger pushed to insert the tag. The needle is removed, and the tag lightly tugged to set the dart. Following tagging, all sharks are released alive and monitored to ensure normal post capture behaviour. Currently, the data is stored with the MWT. Previously data had also been stored with the SSTP.

Results

Sharks tagged in 2025

One training boat trip was undertaken in 2025. The second trip had to be cancelled due to ongoing weather issues. The trip saw the practical training completed for 8 new shark taggers. A total of 123 individuals were tagged, including one spurdog and 122 tope (Figure 2). The majority of the tope tagged were female, 117 individuals in total, while five were male. The spurdog tagged was male, as shown in Figure 2. The angler who caught the spurdog only tagged the first of many that day, they noted that they were caught over a wide area, all of a similar size and mostly male.




Figure 2 The number and sex of small sharks tagged in Manx waters during 2025.

The length of the spurdog was 85.00 cm. Tope ranged in length from 57.00 cm to 167.00 cm, with an average length of 146.01 cm (± 18.71 cm). Figure 3 shows the range and average length of tagged small sharks from 2025 (Appendix 3).




Figure 3 Box plot showing the median length, and interquartile range (i.e., the range in values of the central 50% of the data) of tagged small sharks in 2025. Whiskers indicate the minimum and maximum lengths recorded.

Distribution of sharks tagged in 2025

Sharks were tagged towards the south, southeast and southwest of the island with some also being tagged several miles off Douglas. Notably catches around the Calf of Man were significantly less than the 2024 season. However, this may be a reflection that anglers tend to fish in areas where certain species are known to be found. Nevertheless, in combination with data obtained in subsequent tagging years, this data may contribute to the identification of hotspots, sex aggregations and/or potential nursery areas.

Recaptures

Five Manx recaptures and one Irish were reported in 2025. Originally tagged in Dalby, Niarbyl, and along the west coast in 2022 and 2024. Due to a recording error one tope's tagged location and date is unknown. They were recaptured off Olhao, Portugal, North Spain, Burry Point, South Wales, Southwest of lundy Island and Derbyhaven respectively. The Irish shark was originally tagged in July of 2019 off Greystones, on the east coast of Ireland. The shark was recaptured in August 2025 off Derbyhaven, Isle of Man. This was the longest time between tag and recapture this year, after 2249 days and 130 km from its original tag location.

Since 2013 this brings our known recaptures, both local and non-local, to a total 22. Recaptures provide interesting data, showing tope inhabiting Manx waters are migratory across European waters and as far as the waters of the northwest coast of Africa, over 3000 km from the Island. Highlighting the need to protect them across many jurisdictions and not just Manx waters. This is particularly important in some European areas where the sharks are

landed for market, as has been reported on several occasions with the recapture records. Additionally, these recaptures, substantiated by research findings, suggest there to be a single stock of tope throughout European waters (ICES, 2012).

Comparison of sharks tagged 2013-2025

A total of 809 small sharks have been tagged since 2013 (Table 1). Numbers of tagged individuals have increased over the last four years in comparison to previous years, which were predominantly tope. The 2025 season has seen further improvement of the number of sharks tagged. Such high tagging success is thanks to the continued support of several anglers and angling charters, namely Casey J and Gemini.

Table 1. The number of small sharks per species tagged between 2013 – 2025.

Species	Year												
	2013	2014	2015	2016	2017	2018	2019	2021	2022	2023	2024	2025	
Bull huss	17	1	0	0	0	0	0	0	4	2	0	0	
Spurdog	5	1	1	4	90	14	8	31	12	3	5	1	
Thornback ray	0	0	0	0	0	0	0	0	7	0	0	0	
Tope	28	21	20	12	40	30	10	22	132	58	108	122	
Total	50	23	21	16	130	44	18	53	155	63	113	123	

Spurdog

One male spurdog was tagged during the 2025 tagging effort. The angler who caught and tagged this fish noted that they caught multiple spurdog that day mostly male of a similar size. Females have been more frequently tagged than males across the entire programme, as shown in Figure 4. The 2024 distribution of spurdog shows they were tagged in waters around the south of the Island, the individual caught in 2025 was 3.5 Nm east of Douglas. Spurdog travel in large, dense aggregations, segregated by size and sex (Henderson *et al.*, 2002). This makes it unsurprising that females have been primarily tagged but highlights that Manx waters may be crucial migratory routes or potential pupping grounds for female spurdog.




Figure 4 The number of individual spurdog females (blue) and males (orange), as well as the unidentified individuals (grey) between 2013 – 2025. The two lines denote the break in survey effort in 2020 due to COVID-19.

Figure 5 illustrates the average length of female spurdog between 2013 to 2024 (Appendix 3). The data shows the average to be consistent between 2013 and 2018 before decreasing over the period between 2018 and 2021. 2022 showed a positive increase in the average length of tagged spurdog continuing into the 2023 and 2024 season (Figure 6). It is also important to take the spatial distribution and number of spurdog caught and tagged into account when interpreting this data. The minimum total length for females at sexual maturity (50 % certainty) is 74 cm (Henderson *et al.*, 2002). Throughout the programme, the majority of tagged females were above this threshold and therefore likely to be sexually mature, indicating that Manx waters may provide nursery grounds for this species.




Figure 5 The minimum and maximum length (cm) of tagged female spurdog (whiskers) and average length between 2013 – 2024. Dashed line indicates minimum length of maturity (74 cm). One individual was tagged in 2014, 2015 and 2022 respectively, therefore data was modelled for these years to generate an average length.

Tope

Whilst the sex ratio of tagged tope shows annual variation, females have been more frequently tagged than males over the last six years, as shown in Figure 6. The total number of females caught and tagged has increased in the past years compared to previous years, the number of males caught is both lower and varies annually. The proportion of females to males could be interpreted to suggest that Manx waters may predominantly be used by females, perhaps utilising the area as a small shark nursery ground. However, this is deemed unlikely as the majority of tagged female tope do not meet the 50% certainty of maturity threshold of 155 cm TL (Figure 7; Dureuil, 2013). Male tope have been consistently above the equivalent threshold of 121 cm TL (Figure 9; Dureuil, 2013) until this year. This may be due to this species typically being partially segregated by size and sex (Walker *et al.*, 2008). It's notable that the average length of male tope was smaller than that of previous years.

The distribution of tope caught during the 2025 survey period were around most of the southern half of the Island, particularly off Peel, Douglas and Port St Mary. They were also caught up the west coast and closer to the Calf. The distribution shows significantly less caught around the Calf than the previous year. This could be, in part, a result of the second boat trip not taking place, which would normally target the south of the Island including the Calf.




Figure 6 The number of individual tope females (blue) and males (orange), as well as the unidentified individuals (grey) between 2013 – 2025. The curved dotted line indicates average number of males and females over time.




Figure 7 The average length (cm) of total tagged tope (orange); average length (cm) of female tagged tope (blue); average length (cm) of male tagged tope (green); and average length for sexually mature females (155 cm, dashed blue) and males (121 cm, dashed green) between 2013 – 2025.

Bull huss

Despite some success in previous years, no bull huss were tagged this season. Prior to the 2022 and 2023 tagging effort, bull huss had only been tagged in 2013 and 2014 (Table 1). It is unclear as to why more bull huss have not been tagged throughout the programme, as research suggests bull huss are locally abundant in the Irish Sea and populations are slowly increasing (ICES-WGEF, 2019). This could be a reflection of local fishing pressures in Manx waters.

Thornback ray

No thornback rays were caught this season.

Overview of Programme

The total number of tagged individuals for the 2025 survey period continued to increase on past year's numbers. In 2025, tope were almost exclusively reported and tagged. The majority of individuals caught and tagged were female. Figure 6 shows a slight decrease in size and numbers of male sharks from last year and an increase in numbers of females. One spurdog was caught and tagged this year, with more of similar sex and similar size caught that same day however, the data was not officially recorded. Bull huss and thornback ray were not recorded this year.

The average length of tope and the individual spurdog decreased from the year prior. Average size of small sharks tagged remains variable over the years across both species. Tagging a wider range of shark sizes may provide the programme with more insight into whether Manx waters are important for part of their reproductive cycles, such as for breeding or as a nursery for young.

The south of the island remains a key area for small shark species and may reflect anglers preferences for certain areas. Small sharks have consistently been tagged south of the Calf

of Man, north of Peel, south of Douglas and off Niarbyl. Effort was made in the 2022 survey season to tag small sharks towards the more northerly parts of the Island, to gain an understanding of the sex and size ratios of species in this area. Efforts to maintain a better coverage of the island is important as it helps to remove the bias of fishing. Appendix 4 contains maps containing all tagging locations between 2013 – 2025.

Conclusions and Recommendations

In total, 809 small sharks have been tagged since 2013. An additional 13 small sharks were tagged prior to the formal commencement of the Small Shark Tagging Project in 2013. Increased tagging effort in 2022 and 2023 saw the tagging of bull huss, which had not been recorded as caught or tagged since 2014. For the first time in the programme, the thornback ray was also tagged in 2022 after an effort was made to fish in more northern waters surrounding the island. Thornback ray remained elusive for the following seasons however this could be due to a number of reasons including less targeted fishing, reduced fishing trips, fishing trips taking place in a less diverse range of locations as well as other environmental factors. Gaining more data on a greater variety of small shark species is hugely beneficial in furthering our understanding of where and how these species use Manx waters. This knowledge can help us to further maintain and protect our fisheries surrounding the island.

The project has recorded six recaptures this year, five of which have occurred from elasmobranchs originally tagged on the Island and one recapture from Ireland. Further highlighting the need to protect these species across many jurisdictions, not just the Isle of Man. To help to increase the probability of successful recapture, it is proposed that the continuation of tagging workshops will be key to increase the number of sharks tagged in our waters. Whilst this method can be considered as invasive for the species involved, it is a good way to minimise the disruption to the sharks and rays as the fishers are able to carry out the research whilst they are already fishing. In the same way, it helps to reduce the monetary and environmental cost of separate tagging efforts. Involving fishers and educating them is also an imperative part to achieving healthier waters around our island. Overall, it is too soon to determine whether small sharks are utilising Manx waters for migrations, feed grounds or as part of their reproductive cycles. Continued tagging and recapture of previously tagged individuals are crucial to obtain useful information about the distribution and population structure of small sharks in Manx waters.

Currently, small sharks have been tagged in several Marine Nature Reserves (MNRs) including West Coast, Calf and Wart Bank, Baie ny Carrickey, Langness, Little Ness and Ramsey Bay. These sites only cover up to the 3 nm boundary of Manx waters and are not formally designated to protect small shark species. This is in part due to small shark species not receiving formal protection in the Isle of Man currently. However, small sharks will benefit from these MNRs due to restrictions against damaging fishing methods.

Based on our current understanding of the tagged shark species, we recommend greater protection in the form of restrictions or reserve formation/extension into the 3-12 nm zone of the Calf and Wart Bank, Langness and West Coast MNRs. Extending protection into the wider 3-12 nm zone around the Island is crucial to protect these areas from damaging marine developments and fishing.

Manx Wildlife Trust is grateful for the support of this programme from the angling community, including the charter boats Casey J and Gemini, and is optimistic for future data collection.

References

- Baino, R., Serena, F., Ragonese, S., Rey, J. and Rinelli, P. 2001. Catch composition and abundance of Elasmobranchs based on the MEDITS program. *Rapp. Comm. Int. Mer Médit.* **36**, 234.
- Bovcon, N.D., Cochia, P.D., Navoa, X., Ledesma, P., Caille, G.M. and Baigun, C.R.M. 2018. First report on a pupping area of the tope shark *Galeorhinus galeus* (Carcharhiniformes, Triakidae) in the south-west Atlantic. *Journal of Fish Biology.* **93**(6), 1229-1232.
- Bubley, W.J., Sulikowski, J.A., Koester, D.M. and Tsang, P.C. 2013. Using a multi-parameter approach to reassess maturity of spiny dogfish, *Squalus acanthias*, following increased fishing pressure in the Western North Atlantic. *Fisheries Research.* **147**, 202-212.
- Capapé, C., Vergne, Y., Vianet, R., Guélorget, O. and Quignard, J. 2006. Biological observations on the nursehound, *Scyliorhinus stellaris* (Linnaeus, 1758) (Chondrichthyes: Scyliorhinidae) in captivity. *Acta Adriatica.* **47**(1), 29-36.
- Castro, J.I. 1983. *The sharks of North American waters.* Texas A & M University Press, College Station.
- Compagno, L.J.V. 1984. *FAO species catalogue. Vol. 4. Sharks of the world. An annotated and illustrated catalogue of shark species known to date.* FAO Fisheries Synopsis No. 125, Volume 4, Part 1.
- Daan, N., Heessen, H.J.L. and ter Hofstede, R. 2005. North Sea Elasmobranchs: distribution, abundance and biodiversity. ICES CM 2005/N:06.
- Dureuil, M. 2013. Status and conservation of sharks in the Northeast Atlantic. M.Sc. Thesis. Biological Oceanography. University of Kiel, Kiel.
- Ebert, D.A. 2003. *Sharks, Rays and Chimaeras of California.*
- Ebert, D.A., Fowler, S. and Compagno, L. 2013. *Sharks of the World. A Fully Illustrated Guide.* Wild Nature Press, Plymouth, United Kingdom.
- Ellis, J. 2016. *Raja clavata.* *The IUCN Red List of Threatened Species 2016:* e.T39399A103110667. [Online]. Available at: <https://dx.doi.org/10.2305/IUCN.UK.2016-3.RLTS.T39399A103110667.en>. Accessed 26.01.23.
- Ellis, J., Dulvy, N., Walls, R. and Serena, F. 2016. *Raja clavata (Europe assessment).* *The IUCN Red List of Threatened Species 2016:* e.T39399A103111648 [Online]. Available at: <https://www.iucnredlist.org/species/39399/103111648>. Accessed on 04.02.23.
- Ellis, J., Serena, F., Mancusi, C., Haka, F., Morey, G., Guallart, J. and Schembri, T. 2015a. *Scyliorhinus stellaris (Europe assessment).* *The IUCN Red List of Threatened Species 2015:* e.T161484A48923567 [Online]. Accessed on 27.01.23.
- Ellis, J., Soldo, A., Dureuil, M. and Fordham, S. 2015b. *Squalus acanthias (Europe assessment).* *The IUCN Red List of Threatened Species 2015:* e.T91209505A48910866 [Online]. Accessed on 27.01.23.
- Finucci, B., Derrick, D. and Pacoureaux, N. 2021. *Scyliorhinus stellaris.* *The IUCN Red List of Threatened Species 2021:* e.T161484A124493465 [Online]. Available at:

<https://dx.doi.org/10.2305/IUCN.UK.2021-2.RLTS.T161484A124493465.en>.
Accessed on 26.01.23.

- Fordham, S., Fowler, S.L., Coelho, R.P., Goldman, K. and Francis, M.P. 2016. *Squalus acanthias*. *The IUCN Red List of Threatened Species* 2016: e.T91209505A2898271 [Online]. Available at: <https://www.iucnredlist.org/species/91209505/2898271>. Accessed 30.06.20.
- Francis, M.P. and Mulligan, K.P. 1998. Age and growth of New Zealand school shark, *Galeorhinus galeus*. *New Zealand Journal of Marine and Freshwater Research*. **32**(3), 427-440.
- Henderson, A.C., Flannery, K. and Dunne, J. 2002. Growth and Reproduction in Spiny Dogfish *Squalus acanthias* L. (Elasmobranchii: Squalidae), from the West Coast of Ireland. *Sarsia*. **87**(5), 350-361.
- Holden, M.J. 1975. The fecundity of *Raja clavata* in British waters. *Journal du Conseil International de l'Exploration de la Mer*. **36**, 110-118.
- Holden, M.J. and Horrod, R.G. 1979. The migration of tope, *Galeorhinus galeus* (L), in the eastern North Atlantic as determined by tagging. *ICES Journal of Marine Science*. **38**, 314-317.
- ICES. 2011. Report of the Working Group on Elasmobranch Fishes (WGEF), Copenhagen, Denmark, 20-24 June 2011. ICES CM 2011/ACOM:19. International Council for the Exploration of the Sea (ICES), Denmark.
- ICES. 2012. Report of the Working Group on Elasmobranch Fishes (WGEF), Lisbon, Portugal, 19-26 June 2007. ICES CM 2012/ACOM:19. International Council for the Exploration of the Sea (ICES), Denmark.
- ICES-WGEF. 2019. Working Group on Elasmobranch Fishes (WGEF). ICES Scientific Reports 1:25. International Council for the Exploration of the Seas, Copenhagen, Denmark.
- Jenkins, J.T. 1958. *The Fishes of the British Isles*. London: Frederick Warne & Co.
- Jones, T.S. and Ugland, K.I. 2001. Reproduction of female spiny dogfish, *Squalus acanthias*, in the Oslofjord. *Fishery Bulletin*. **99**, 685-690.
- Kadri, H., Marouani, S., Saidi, B., Bradai, M.N., Bouain, A. and Eric, M. 2014. Age, growth, sexual maturity and reproduction of the thornback ray, *Raja clavata* (L.) of the Gulf of Gabe`s (south-central Mediterranean Sea). *Marine Biology Research*. **10**, 416-425.
- McCully, S.R., Dureuil, M. and Farrell, E. 2015. *Galeorhinus galeus* (Europe assessment). *The IUCN Red List of Threatened Species* 2015: e.T39352A48938136 [Online]. Available at: <https://www.iucnredlist.org/species/39352/48938136>. Accessed on 03.02.23.
- McCully, S.R., Scott, F. and Ellis, J.R. 2012. Lengths at maturity and conversion factors for skates (Rajidae) around the British Isles, with an analysis of data in the literature. *ICES Journal of Marine Science*. **69**, 1812-1822.
- McHugh, M., Sims, D.W., Partridge, J.C. and Genner, M.J. 2011. A century later: Long-term change of an inshore temperate marine fish assemblage. *Journal of Sea Research*. **65**, 187-194.

- McMillan, M.N., Huvneers, C., Semmens, J.M., Gillanders, B.M. and Durif, H.e.C. 2018. Partial female migration and cool-water migration pathways in an overfished shark. *ICES Journal of Marine Science*. **76**(4),1083-1093.
- Olsen, A.M. 1954. The biology, migration, and growth rate of the school shark, *Galeorhinus australis* (Macleay) (Carcharhinidae) in south-eastern Australian waters. *Australian Journal of Marine and Freshwater Research*. **5**, 353-410.
- Rodriguez-Cabello, C., Sánchez, F. and Velasco, F. 2005. Growth of lesser spotted dogfish (*Scyliorhinus canicula* L., 1758) in the Cantabrian Sea, based on tag-recapture data. *Journal of Northwest Atlantic Fishery Science*. **37**, 131-140.
- Ryland, J.S. and Ajayi, T.O. 1984. Growth and population dynamics of three *Raja* species in Carmarthen Bay, British Isles. *Journal de Conseil Internationale de Exploration de la Mer*. **41**, 111-120.
- Serra-Pereira, B., Figueiredo, I. and Serrano Gordo, L. 2011. Maturation, fecundity, and spawning strategy of the thornback ray, *Raja clavata*: do reproductive characteristics vary regionally. *Marine Biology*. **158**(2), 187-2197.
- Sherley, R.B., Winker, H., Rigby, C.L., Kyne, P.M., Pollom, R., Pacoureau, N., Herman, K., Carlson, J.K., Yin, J.S., Kindsvater, H.K. and Dulvy, N.K. 2020. Estimating IUCN Red List population reduction: JARA—A decision-support tool applied to pelagic sharks. *Conservation Letters*. **13**(2), e12688.
- Veríssimo, A., McDowell, J.R. and Graves, J.E. 2010. Global population structure of the spiny dogfish, *Squalus acanthias*, a temperate shark with an antitropical distribution. *Molecular Ecology*. **19**, 1651-1662.
- Walker, P.A. 1998. *Fleeting Images: Dynamics of North Sea Ray Populations*. Ph.D. Thesis, University of Amsterdam.
- Walker, T.I. 1999. Chapter 24. *Galeorhinus galeus* fisheries of the world. In: Shotton, R. (ed.), *Case studies of management of elasmobranch fisheries*. *FAO Fisheries Technical Paper 378/2*, pp. 728-773. Food and Agriculture Organization of the United Nations, Rome, Italy.
- Walker, T.I. 2005. Chapter 4 *Reproduction in fisheries science*. In: W. C. Hamlett (ed.), *Reproductive biology and phylogeny of Chondrichthyes: sharks, batoids, and chimaeras*, pp. 81-127. Science Publishers, Inc., Enfield, New Hampshire 03478, USA.
- Walker, T.I., Cavanagh, R.D., Stevens, J.D., Carlisle, A.B., Chiaramonte, G.E., Domingo, A., Ebert, D.A., Mancusi, C.M., Massa, A., McCord, M., Morey, G., Paul, L.J., Serena, F. and Vooren, C.M. 2006. *Galeorhinus galeus*. *The IUCN Red List of Threatened Species* 2006: e.T39352A10212764 [Online]. Available at: <http://dx.doi.org/10.2305/IUCN.UK.2006.RLTS.T39352A10212764.en>. Accessed 30.06.20.
- Walker, T.I., Rigby, C.L., Pacoureau, N., Ellis, J., Kulka, D.W., Chiaramonte, G.E. and Herman, K. 2020. *Galeorhinus galeus*. *The IUCN Red List of Threatened Species* 2020: e.T39352A2907336 [Online]. Available at: <https://dx.doi.org/10.2305/IUCN.UK.2020-2.RLTS.T39352A2907336.en>. Accessed on 26.01.23.
- Walker, T.I., Taylor, B.L., Brown, L.P. and Punt, A.E. 2008. Chapter 32. Embracing movement and stock structure for assessment of *Galeorhinus galeus* harvested off southern

- Australia. In: Camhi, M.D., Pikitch, E.K. and Babcock, E.A. (eds), *Sharks of the Open Ocean: Biology, Fisheries and Conservation*, pp. 369–392. Blackwell Publishing, Oxford, United Kingdom.
- Wiegand, J., Hunter, E. and Dulvy, N.K. 2011. Are spatial closures better than size limits for halting the decline of the North Sea thornback ray, *Raja clavata*? *Marine and Freshwater Research*. **62**, 722-733.
- Winker, H, Pacoureaux, N. and Sherley, R.B. 2020. JARA: 'Just Another Red List Assessment'. *BioRxiv Preprint*: <http://dx.doi.org/10.1101/672899>.

Appendices

Appendix 1. Tagging guidance crib sheet.

SSTP Micro-Tag Guidelines



Micro-tags are inserted using a tagging gun. When not in use we recommend that you use the needle guard to prevent injury!

1. Fit a strip of micro-tags into the tagging gun
2. Insert the needle at an angle of 45°
3. Push the trigger to insert a single tag into the fish
4. Remove the needle and give the tag a short tug to set the barb

Submit your data to Eleanor – eleanor@manxwt.org.uk or drop in at 7-8 Market Place, Peel, IM5 1XF or online at: www.tagsharks.com

These tagging guidelines are for tagging guns and micro-tags ONLY. SSTP minimum sizes are set to protect fish, if for any reason you are unsure about tagging please **do not tag the fish!**

SSTP Micro-Tag Minimum Sizes	
Common Skate (wingspan)	Canula
Tope	65cm (2.8lb)
Spurdog	65cm (2.3lb)
Smooth-hound	70cm (2.2lb)
Bull Huss	65cm (2.5lb)
Rays (wingspan)	35cm (2.1lb)

Appendix 2. Record card.

Please send details to Eleanor by email: eleanor@manxwt.org.uk
Or drop in/post to: 7-8 Market Place, Peel, IM5 1XF


Name/s: _____

Email address: _____




Date: _____ Time start: _____ Time end _____

Location (please circle): NE NW SW SE

Lat/Long (this will NOT be made public):
_____ N _____ W



In association with:

Tag No.	Species	Sex	Length (cm)	Girth (cm)	Condition

Appendix 3. The range and average length (cm) of tagged sharks between 2013 – 2025.

Table 1. The range and average length (cm) of tagged bull huss for 2022. N.B.: averages for 2014 were not able to be calculated as only one individual was tagged.

Year	Length range (cm)		Average length (cm)	Standard deviation (±)
	Minimum	Maximum		
2013	63.00	110.00	91.06	13.71
2014	89.00		-	-
2022	74.00	100.00	86.00	9.90
2023	96.00	107.00	101.50	7.78

Table 2. The range and average length (cm) of tagged spurdog between 2013 – 2025. N.B.: averages for 2014, 2015, 2022 and 2025 were not able to be calculated as only one individual was tagged in each of these years respectively.

Year	Length range (cm)		Average length (cm)	Standard deviation (±)
	Minimum	Maximum		
2013	75.00	107.00	94.17	11.92
2014	100.00		-	-
2015	101.00		-	-
2016	101.00	108.00	104.00	3.16
2017	74.50	113.00	100.49	6.76
2018	98.00	110.00	104.93	3.67
2019	90.00	102.00	96.75	3.83
2021	50.00	115.00	84.48	19.02
2022	70.00	105.00	96.83	9.43
2023	80.00	94.00	86.00	7.21
2024	75.00	110.00	93.00	18.06
2025	85.00		-	-

Table 3. The range and average length (cm) of tagged thornback ray for 2022.

Year	Length range (cm)		Average length (cm)	Standard deviation (±)
	Minimum	Maximum		
2022	60.00	145.00	103.14	36.14

Table 4. The range and average length (cm) of tagged tope between 2013 – 2025.

Year	Length range (cm)		Average length (cm)	Standard deviation (±)
	Minimum	Maximum		
2013	110.00	156.00	141.71	12.32
2014	94.00	145.00	124.95	14.95
2015	80.00	153.00	122.00	24.10
2016	89.00	157.00	130.70	24.07
2017	99.00	168.00	142.38	14.51
2018	78.00	171.00	145.93	21.97
2019	110.00	167.00	148.10	17.88
2021	60.00	168.00	127.91	26.54
2022	99.00	173.00	149.24	14.03
2023	105.50	165.10	145.13	13.90
2024	63.60	171.00	149.64	12.90
2025	57.00	167.00	146.01	18.71

Appendix 4. The capture location of small sharks tagged in Manx waters during 2013 (a), 2014 (b), 2015 (c), 2016 (d), 2017 (e), 2018 (f), 2019 (g), 2021 (h), 2022 (i), 2023(j), 2024(k) and 2025(l).

